

## ROTATION DAY 11: ANGULAR MOMENTUM

Rigid body about a fixed axis:  $L = I\omega$     If  $\sum \tau_{\text{ext}} = 0$ :  $L_i = L_f$

Later today: a moving particle can also have angular momentum about a point, with  $L = mvr_{\perp}$ .

### CONSERVATION OF ANGULAR MOMENTUM

#### WE DO    The Skater Spin

A figure skater has  $I = 4.5 \text{ kg}\cdot\text{m}^2$  with arms extended and spins at 1.5 rev/s. She pulls her arms in, reducing  $I$  to 1.8  $\text{kg}\cdot\text{m}^2$ .

(a) Convert  $\omega_i$  to rad/s.

(c) Find  $\omega_f$ .

(b) Is there an external torque?  $\rightarrow L$  is conserved.

(d) Convert  $\omega_f$  to rev/s. How many times faster?

(e) Did her kinetic energy change? Calculate  $K_i$  and  $K_f$ . Where did the extra energy come from?

### COLLISIONS ON A TURNTABLE

#### WE DO    Merry-Go-Round + Person

A 150 kg merry-go-round (solid disk, radius 2.0 m) spins at 0.50 rad/s. A 60 kg person jumps onto the edge from rest (radially).

(a) Find  $I_{\text{disk}} = \frac{1}{2}MR^2 =$  \_\_\_\_\_

(d) Apply conservation:  $I_i\omega_i = I_f\omega_f$ . Find  $\omega_f$ .

(b) Find  $I_{\text{person}} = mR^2$  (treat as point mass) = \_\_\_\_\_

(e) Is kinetic energy conserved? Calculate both. This is like an inelastic collision!

(c)  $I_{\text{total after}} = I_{\text{disk}} + I_{\text{person}} =$  \_\_\_\_\_



#### YOU DO    Catch on a Turntable

A person stands at the center of a stationary turntable ( $I_{\text{system}} = 5.0 \text{ kg}\cdot\text{m}^2$ ). A friend throws a ball (mass 0.50 kg) horizontally. The ball lands in the person's hands at  $r = 0.80 \text{ m}$  from the axis, moving at 10 m/s tangentially.

(a) What is the ball's angular momentum about the axis before being caught? (Hint:  $L = mvr$  for tangential motion.)


(b) After the catch, what is the total  $I$  of the system (turntable + person + ball)?

(c) Find  $\omega$  after the catch.

## ANGULAR MOMENTUM IS A VECTOR!

### DIRECTION OF $\vec{L}$

So far we've treated  $L$  as a number — positive or negative. But angular momentum is a **vector**. Its direction is along the **axis of rotation**, determined by the **right-hand rule**: curl your fingers in the direction of spin, and your thumb points in the direction of  $\vec{L}$ .




**Key insight:** Since  $\vec{L}$  is a vector, changing its *direction* — not just its magnitude — requires a net external torque. The relationship  $\vec{\tau}_{\text{net}} = \frac{\Delta \vec{L}}{\Delta t}$  tells us that torque applied over time changes angular momentum. **This takes torque  $\times$  time, not work!**

## DEMO: THE BIKE WHEEL GYROSCOPE

### DEMO Spinning Bike Wheel on a String

Your teacher holds a spinning bike wheel by one end of its axle and releases the other end. Instead of tipping straight down right away, the wheel begins to rotate (precess) horizontally.



**What you observe:**

(a) Which direction does  $\vec{L}$  point when the wheel spins? (Use the right-hand rule.)

(b) Gravity pulls the free end down. What direction is the torque  $\vec{\tau} = \vec{r} \times \vec{F}$ ? (Hint: it's perpendicular to both the axle and gravity — it's *horizontal*.)

(c) Since  $\Delta \vec{L} = \vec{\tau} \Delta t$  is horizontal, does  $\vec{L}$  tilt *down* or rotate *sideways*?

**Why doesn't it immediately tip straight down?** In this idealized model, the torque from gravity is *perpendicular* to  $\vec{L}$ , so it mainly changes the *direction* of  $\vec{L}$  rather than its magnitude — just like a centripetal force changes the direction of velocity without changing speed. The wheel turns sideways (precesses) instead of just dropping straight down right away.

### THINK Predictions


(d) If you spin the wheel *faster*, the precession rate: increases / decreases / stays the same

**L OF A POINT MASS IN A STRAIGHT LINE**

The same angular momentum idea also works when an object is *not* spinning as a rigid body. A particle doesn't have to move in a circle to have angular momentum about a point! For any particle with momentum  $p = mv$ :

$$L = mvr_{\perp}$$

where  $r_{\perp}$  is the **perpendicular distance** from the axis to the particle's line of motion (the "closest approach distance").



**Equivalently:**  $L = mvr \sin \theta$ , where  $\theta$  is the angle between  $\vec{r}$  and  $\vec{v}$ . When motion is tangential,  $\sin \theta = 1$ . When the particle moves straight toward or away from the axis,  $\sin \theta = 0$  and  $L = 0$ .

**Big picture:** An object's total angular momentum = *orbital*  $L$  (motion of its center of mass about the axis) + *spin*  $L$  (rotation about its own center). Think: Earth orbits the Sun **and** spins on its axis.

**YOU DO Ball Past a Pivot**

A 0.50 kg ball travels at 8.0 m/s in a straight line that passes 1.2 m from point O.

(a)  $L = mvr_{\perp} = \underline{\hspace{2cm}}$  kg·m<sup>2</sup>/s (b) Does L change along the path (no external torques)?

(c) Using the right-hand rule, is  $\vec{L}$  into or out of the page? \_\_\_\_\_

**ENGINEER The Helicopter Problem**

The engine applies a torque to spin the main rotor blades **clockwise** (as seen from above).

(a) By Newton's 3rd Law, what must happen to the *body* of the helicopter?

(b) The small tail rotor pushes air sideways. Explain its purpose using the concept of torque.

**ANGULAR IMPULSE****ANGULAR IMPULSE–MOMENTUM THEOREM**

Just as  $F\Delta t = \Delta p$ :  $\tau\Delta t = \Delta L = I_f\omega_f - I_i\omega_i$  — a torque applied for a time changes angular momentum.

**QUICK Conceptual Checks** True/False. Correct any false statements.

**(a)** If net external torque is zero, angular velocity must be constant. \_\_\_\_ **(b)** A figure skater's angular momentum increases when she pulls her arms in. \_\_\_\_

**(c)** Angular momentum is always conserved. \_\_\_\_ **(d)** A ball moving in a straight line can have angular momentum about a point not on the line. \_\_\_\_

**SPORT The High Dive**

A diver leaves the board with angular momentum  $L$ . Air resistance is negligible.

**Tuck:** She grabs her knees.  $I$  decreases.

**Layout:** She straightens out.  $I$  increases.

$\omega$ : increases / decreases / stays the same

$\omega$ : increases / decreases / stays the same

$L$ : increases / decreases / stays the same

$L$ : increases / decreases / stays the same

**Key check:** Did  $L$  change during the airtime? \_\_\_\_ Why or why not?

**YOU DO Spinning Platform with Weights**

A person stands on a frictionless turntable holding two 3.0 kg masses at arm's length (0.80 m from axis). The system spins at 2.0 rad/s. Total  $I_{\text{person+turntable}}$  (without masses) = 4.0 kg·m<sup>2</sup>.

**(a)**  $I_{\text{total initial}}$  (include two masses at 0.80 m):

**(c)** Find the new  $\omega$ .

**(b)** Masses pulled to 0.20 m.  $I_{\text{total final}}$ :

**(d)** By what factor did rotational KE change?

## EXIT TICKET & HOMEWORK

### EXIT TICKET

A satellite has a 200 kg main body (model as a uniform sphere,  $R = 0.50$  m) spinning at 2.0 rad/s about its own center of mass. It extends two solar panels (total mass 20 kg, modeled as point masses at 3.0 m from the center). What is the new spin rate?

### HOMEWORK

#### 1. Basic Conservation

A disk ( $I = 0.40$  kg·m<sup>2</sup>) spinning at 8.0 rad/s is dropped onto an identical stationary disk (same  $I$ ). They stick together. Find  $\omega_f$ .

#### 2. Neutron Star

When a massive star collapses, its core shrinks from  $R \approx 700,000$  km to  $R \approx 10$  km while approximately conserving angular momentum. If the original star rotated once every 30 days, estimate the rotation period of the neutron star. (Assume uniform spheres —  $I = \frac{2}{5}MR^2$ .)

#### 3. Angular Impulse

A torque of 12 N·m is applied to a wheel ( $I = 3.0$  kg·m<sup>2</sup>) for 4.0 s. The wheel starts from rest. (a) Find  $\Delta L$ . (b) Find  $\omega_f$ .

#### 4. AP Comparison

Compare and contrast conservation of linear momentum ( $p = mv$ ) and conservation of angular momentum ( $L = I\omega$ ). When is each conserved? Can one be conserved while the other isn't? Give a specific example.

**Next Class:** Tomorrow is full rotation review. Bring everything — kinematics through angular momentum. The unit test is Day 13.